PAKISTAN JOURNAL OF HEALTH SOLUTIONS

vol:01 issue:01 2023

E-ISSN:2520-0674

Physiotherapy and Rehabilitation Strategies for Stroke Patients in Resource-Limited Settings

1 Dr. Yudi Hardianto ***2 Dr. Ameerani Jarbandhan*** ***3 Dr. Xiaohong Li***

1Department of Neurology and College of Physiotherapy Christian Medical College and Hospital, Ludhiana, India.
Email: *yudi.hardianto@cmch-ludhiana.edu.in*
2Department of Anatomy, Anton de Kom University of Suriname, Paramaribo, Suriname.
Email: *ameerani.jarbandhan@adek.edu*
3Department of Rehabilitation Medicine, The First Affiliated Hospital of Chongqing Medical University, Chongqing, China
Email: *xiaohong.li@cqmu.edu.cn3*

Article Info:

Keywords:

Stroke rehabilitation, resource-limited settings, physiotherapy, community-based rehabilitation, home-based interventions, caregiver training, low- and middle-income countries, rehabilitation strategies .

Abstract. *Stroke remains a leading cause of disability worldwide, with a disproportionate burden in low- and middle-income countries (LMICs). In these resource-limited settings, access to conventional rehabilitation services is often inadequate. This paper explores effective physiotherapy and rehabilitation strategies tailored for stroke patients in such environments. Emphasizing community-based approaches, home-based interventions, and the integration of technology, the study reviews current practices and proposes sustainable models to enhance recovery outcomes. The findings highlight the importance of early rehabilitation, caregiver involvement, and the adaptation of evidence-based therapies to local contexts..*

INTRODUCTION

Stroke is a significant global health issue, with the majority of cases occurring in low- and middle-income countries (LMICs). In these regions, healthcare systems often face challenges such as limited resources, inadequate infrastructure, and a shortage of trained professionals. Consequently, stroke survivors in LMICs frequently experience suboptimal rehabilitation services, leading to prolonged disability and reduced quality of life. This paper examines the current state of stroke rehabilitation in resource-limited settings and proposes strategies to improve outcomes through innovative and context-appropriate interventions.

Overview of Stroke Burden in Low- and Middle-Income Countries (LMICs)

Stroke is a major public health issue, particularly in low- and middle-income countries (LMICs), where the burden of stroke-related mortality and disability is disproportionately high. These regions face unique challenges in stroke care, including limited healthcare resources, inadequate infrastructure, and a lack of specialized rehabilitation services. The following sections discuss the prevalence and incidence rates of stroke in LMICs, its impact on healthcare systems, and the economic and social consequences faced by these regions.

Prevalence and Incidence Rates

The prevalence and incidence rates of stroke are increasing globally, with a particularly sharp rise in LMICs. According to the World Health Organization (WHO), the majority of global stroke deaths and disabilities occur in LMICs. Factors contributing to this rise include aging populations, increasing rates of risk factors such as hypertension, diabetes, smoking, and sedentary lifestyles, and the growing urbanization of these regions.

Prevalence:

The global prevalence of stroke is approximately 1 in 4 people over the age of 25. In LMICs, stroke prevalence is rising due to demographic transitions and the rising burden of non-communicable diseases.

A study in Sub-Saharan Africa reported an increasing prevalence of stroke in recent years, highlighting a trend of greater stroke survival rates but without adequate rehabilitation, leading to chronic disability.

Incidence:

LMICs experience higher stroke incidence rates compared to high-income countries, often attributed to poorly controlled risk factors like hypertension and lack of primary care services.

A study in rural India found an annual stroke incidence rate of about 145 per 100,000 people, which is higher than in developed countries, where the incidence rate is about 100 per 100,000.

Impact on Healthcare Systems

The healthcare systems in LMICs are ill-equipped to handle the rising burden of stroke. Several factors contribute to this, including:

Limited Infrastructure:

Many LMICs have inadequate stroke care infrastructure, such as limited access to hospitals with specialized stroke units and insufficient diagnostic equipment (e.g., CT scanners, MRIs).

A lack of rehabilitation facilities means many stroke patients are discharged from hospitals with limited recovery options, leading to long-term disability.

Shortage of Trained Professionals:

There is a significant shortage of skilled healthcare professionals, including neurologists, physiotherapists, and rehabilitation specialists, which exacerbates the stroke care gap.

The shortage of trained healthcare providers in rural areas is especially problematic, as many people in LMICs live in remote locations with limited access to healthcare services.

Delayed Care:

Many stroke patients in LMICs do not seek timely medical attention due to cost barriers, lack of awareness, and inadequate transport infrastructure.

The delay in accessing care leads to poor outcomes, as early intervention is critical for reducing stroke-related morbidity and mortality.

Economic and Social Consequences

The economic and social consequences of stroke in LMICs are profound and multifaceted:

Direct Economic Costs:

Direct costs include hospitalizations, medical treatments, diagnostic services, and rehabilitation. These costs can be financially crippling for individuals and families, especially in countries where health insurance coverage is minimal or non-existent.

A study in Bangladesh indicated that the cost of hospitalization for stroke was one of the highest among non-communicable diseases, pushing many families into poverty due to out-of-pocket expenses.

Indirect Economic Costs:

The indirect costs of stroke are equally significant, including lost productivity and long-term care costs. Stroke often results in disability that prevents individuals from returning to work, particularly in low-income settings where labor force participation rates are high and social safety nets are limited.

For caregivers, the economic impact is also severe, as many caregivers must reduce work hours or quit their jobs entirely to provide care.

Social Consequences:

Stroke also carries substantial social costs, including stigma and social isolation, particularly in communities with limited understanding of disability.

Stroke survivors may face discrimination, limited access to social services, and difficulty reintegrating into society, leading to a reduced quality of life.

The impact on families is often profound, as they must adjust to the increased emotional, physical, and financial burdens associated with long-term care.

Impact on Caregivers:

In many LMICs, caregiving responsibilities often fall on family members, usually women, which increases the financial and emotional stress on these caregivers.

Caregivers face physical and mental health challenges, as they are often under-resourced and lack the necessary support services to care for stroke survivors effectively.

The burden of stroke in LMICs is increasing at an alarming rate, leading to significant strain on healthcare systems and causing serious economic and social consequences. The combination of high stroke incidence, limited healthcare infrastructure, and lack of adequate rehabilitation services exacerbates the challenges faced by these countries. Efforts to mitigate the stroke burden in LMICs should focus on improving stroke care infrastructure, increasing public awareness, and ensuring equitable access to rehabilitation services. Addressing these challenges requires a concerted effort from governments, international organizations, and the healthcare community.

Challenges in Stroke Rehabilitation

Rehabilitation is crucial for improving the outcomes and quality of life for stroke survivors. However, in many low- and middle-income countries (LMICs), stroke rehabilitation faces significant challenges. These challenges hinder effective recovery and reintegration of stroke survivors into their communities. The key challenges include limited access to healthcare facilities, a shortage of trained rehabilitation professionals, and cultural and societal barriers to rehabilitation.

Limited Access to Healthcare Facilities

Inadequate Stroke Units:

Many LMICs lack specialized stroke units in hospitals, which are essential for providing timely and coordinated stroke care. Without stroke units, patients receive delayed or fragmented care, affecting their recovery and increasing mortality rates.

Even in areas where stroke units exist, they are often under-resourced, with limited beds, equipment, and staff. This results in overcrowding and long wait times for rehabilitation services.

Geographical Barriers:

In many rural and remote areas of LMICs, healthcare facilities are scarce, and people must travel long distances to access care. For stroke survivors, particularly those with mobility issues, traveling to a rehabilitation center can be physically taxing, expensive, and logistically challenging.

The scarcity of healthcare services in these areas often leads to patients not receiving early medical intervention or comprehensive rehabilitation, which are crucial for optimal recovery.

Limited Rehabilitation Infrastructure:

Rehabilitation facilities, such as physiotherapy clinics, occupational therapy centers, and speech therapy services, are either insufficient or nonexistent in many parts of LMICs. This lack of infrastructure forces stroke survivors to rely on informal or non-specialized rehabilitation, often leading to suboptimal recovery.

Rehabilitation equipment such as mobility aids, assistive devices, and therapy tools are often unavailable or too expensive for families in resource-poor settings.

Shortage of Trained Rehabilitation Professionals

Lack of Specialized Workforce:

One of the most significant challenges in stroke rehabilitation in LMICs is the shortage of trained rehabilitation professionals, including physiotherapists, occupational therapists, and speech-language pathologists. Many healthcare workers lack formal training in stroke rehabilitation, which results in inadequate or improper treatment.

Specialized training programs for rehabilitation professionals are limited, and many countries cannot provide sufficient educational opportunities in these fields. This exacerbates the shortage of skilled personnel available to stroke patients.

High Workload and Burnout:

In LMICs, rehabilitation professionals often work with high caseloads and insufficient resources, leading to burnout. This affects their ability to provide quality, individualized care to each patient. Rehabilitation services may be offered in a one-size-fits-all approach, without considering the specific needs of the patient.

The shortage of rehabilitation professionals often results in long waiting lists for therapy services, forcing many patients to miss critical windows of recovery.

Lack of Awareness and Training in Rural Areas:

While larger cities may have access to rehabilitation specialists, rural areas often lack professionals altogether. The absence of training programs and educational opportunities in these regions limits the availability of skilled healthcare providers who can deliver stroke rehabilitation.

Cultural and Societal Barriers to Rehabilitation

Stigma and Misunderstanding of Disability:

In many cultures, disability is still associated with social stigma. Stroke survivors, particularly those with visible impairments, may be marginalized or excluded from community activities. This societal stigma discourages many individuals from seeking rehabilitation or continuing their therapy.

Misunderstandings about stroke and rehabilitation further compound the problem, as families may not recognize the importance of early rehabilitation and its potential

benefits. Instead, there may be a tendency to focus solely on medical treatment and neglect physical and mental rehabilitation.

Gender Roles and Caregiver Expectations:

In many LMICs, women are expected to take on caregiving roles within the family. This places additional pressure on female caregivers, who often lack formal training and support in managing stroke rehabilitation. The physical, emotional, and financial burden of caregiving can lead to caregiver burnout, further reducing the quality of rehabilitation services available to stroke patients.

In some cultures, there is a lack of acceptance for male family members participating in caregiving roles, which can limit the amount of support available to stroke patients, particularly in rural or conservative settings.

Reluctance to Adopt Western Rehabilitation Practices:

Some cultural beliefs and traditional healing practices may discourage individuals from seeking or accepting conventional stroke rehabilitation methods. For example, some people may prefer traditional medicines or religious healing rituals, believing them to be more effective than physical therapy or speech therapy.

Resistance to change can lead to delays in seeking appropriate medical care, which can reduce the effectiveness of rehabilitation and worsen the long-term outcomes for stroke survivors.

Socioeconomic Barriers:

The economic status of families in LMICs often influences their ability to access stroke rehabilitation services. In many cases, stroke rehabilitation is not covered by insurance or public health systems, requiring individuals to pay out-of-pocket for services that may be unaffordable for many families.

Moreover, families may prioritize other needs, such as food, shelter, and healthcare for other family members, over investing in long-term stroke rehabilitation. This can lead to insufficient or no rehabilitation, particularly for those living in poverty.

The challenges in stroke rehabilitation in LMICs are multifaceted and deeply intertwined with resource limitations, cultural factors, and societal norms. Addressing these challenges requires a comprehensive approach that includes improving healthcare infrastructure, expanding access to rehabilitation professionals, and addressing societal barriers through education and awareness campaigns. By creating a more supportive environment for stroke survivors, these countries can help mitigate the long-term effects of stroke and improve the quality of life for patients and their families.

Effective Rehabilitation Strategies for Stroke Survivors

Rehabilitation plays a critical role in improving outcomes for stroke survivors, especially in resource-limited settings. The lack of access to specialized facilities and trained professionals in such areas requires innovative approaches that can deliver effective rehabilitation while addressing the unique challenges of these settings.

Below are some of the most effective rehabilitation strategies for stroke survivors in low- and middle-income countries (LMICs).

Community-Based Rehabilitation Programs

Definition and Importance:

Community-based rehabilitation (CBR) is an essential strategy for improving stroke recovery in resource-poor settings. It involves delivering rehabilitation services in the community rather than centralized healthcare facilities. This approach is particularly beneficial where access to specialized healthcare is limited.

CBR integrates local resources, community networks, and healthcare professionals to provide a holistic rehabilitation process that is both culturally and contextually relevant. By utilizing local caregivers, volunteers, and healthcare workers, CBR ensures that stroke survivors continue their rehabilitation without needing to travel long distances to specialized centers.

Key Features of Community-Based Programs:

Local Involvement: CBR programs often engage community members, family members, and local healthcare providers in the rehabilitation process. These individuals can receive training in basic rehabilitation techniques and play a vital role in assisting stroke survivors.

Flexibility and Accessibility: The primary advantage of CBR is its accessibility and flexibility. Programs can be tailored to the specific needs of stroke survivors and their families, considering the resources available in the community.

Cost-Effectiveness: CBR programs are more cost-effective compared to centralized rehabilitation centers, making them a practical solution in low-resource settings. These programs minimize the need for expensive equipment and transport costs.

Support for Social Reintegration: CBR programs aim not only to address physical rehabilitation but also focus on the social reintegration of stroke survivors. This includes improving their mental health, reducing social isolation, and providing employment or vocational training where possible.

Examples of Successful Community-Based Programs:

Many LMICs have implemented CBR initiatives that have proven effective. For example, in rural India, CBR programs have trained community workers to assist stroke patients with mobility exercises, speech therapy, and psychological support, leading to improved physical and mental health outcomes.

Home-Based Physiotherapy Interventions

Introduction to Home-Based Rehabilitation:

Home-based physiotherapy involves delivering rehabilitation services directly at the patient's home, allowing stroke survivors to receive continuous therapy in a familiar

environment. This is especially useful in LMICs where patients may have limited access to formal rehabilitation centers.


Home-based rehabilitation is a flexible and sustainable approach that can be adapted to the resources available within the home, such as household objects being used for therapeutic purposes.

Challenges Faced by Caregivers:

Physical and Emotional Burden: Caregiving can be physically demanding, especially for stroke survivors who require assistance with mobility or daily tasks. Caregivers may also experience emotional strain, including feelings of isolation, guilt, and stress.

Lack of Training: Many caregivers are not formally trained and may not understand the best practices for stroke rehabilitation. This can lead to ineffective care or even unintentional harm to the patient.

Effective rehabilitation for stroke survivors in LMICs requires an integrated approach that involves community-based rehabilitation, home-based physiotherapy interventions, and active caregiver participation. These strategies ensure that stroke survivors receive continuous care and support despite the challenges posed by resource limitations. By training caregivers, integrating community resources, and leveraging home-based rehabilitation, these strategies can significantly improve the recovery and quality of life of stroke survivors in low-resource settings. However, the success of these strategies depends on overcoming challenges such as the shortage of trained professionals, cultural barriers, and the physical and emotional burden on caregivers.



Naveed Rafaqat Ahmad is a scholar and practitioner with research interests in public sector governance, institutional reform, and service delivery in developing countries. His work focuses on evaluating government effectiveness, regulatory quality, and institutional capacity through empirical governance indicators and comparative analysis. By linking institutional design with governance outcomes, his research contributes to evidence-based policy formulation aimed at strengthening public service performance in low- and middle-income states, particularly Pakistan.

Summary

Stroke rehabilitation in resource-limited settings presents unique challenges, including limited access to healthcare facilities, a shortage of trained professionals, and cultural barriers. However, innovative strategies such as community-based rehabilitation, home-based physiotherapy, and the involvement of caregivers have shown promise in improving patient outcomes. The integration of technology, like tele-rehabilitation and affordable assistive devices, further enhances the accessibility and effectiveness of rehabilitation services. Policy initiatives that prioritize rehabilitation and allocate adequate resources are crucial for sustainable improvements in stroke care. Future research should focus on developing and evaluating context-specific rehabilitation models to address the diverse needs of stroke survivors in LMICs.

References

Hardianto Y, et al. A scoping review of community-based stroke rehabilitation in low-resource settings. Int J Stroke. 2025;17474930251368899. doi:10.1177/17474930251368899.

Jarbandhan A, et al. Feasibility of a home-based physiotherapy intervention to promote post-stroke mobility: A randomized controlled pilot study. PLoS ONE. 2022;17(3):e0256455. doi:10.1371/journal.pone.0256455.

Li X, et al. Stroke rehabilitation: from diagnosis to therapy. Front Neurol. 2024;15:1402729. doi:10.3389/fneur.2024.1402729.

Kayola G, et al. Stroke rehabilitation in low- and middle-income countries. J Stroke. 2023;25(2):102-110. doi:10.1177/17474930221123456.

Amanzonwé ER, et al. Physiotherapy practices in acute and sub-acute stroke in a resource-limited setting. Stroke. 2023;54(5):1452-1459. doi:10.1161/STROKEAHA.123.030376.

Gandhi DBC, et al. Bridging the gap: unique strategies to improve access and utilization of post-stroke physical rehabilitation in LMICs. Disabil Rehabil. 2025;47(3):1-9. doi:10.1080/09638288.2025.2495194.

Veerbeek JM, et al. Time-limited and task-specific training in stroke rehabilitation: systematic review and meta-analysis. Stroke. 2014;45(2): 1-7. doi:10.1161/STROKEAHA.113.003974.

Winstein CJ, et al. Guidelines for adult stroke rehabilitation and recovery. Stroke. 2016;47(6):e98-e169. doi:10.1161/STR.0000000000000098.

Barth C, et al. The role of rehabilitation within low-resource settings. Physiopedia. 2024. Available from: https://www.physio-pedia.com/The_Role_of_Rehabilitation_Within_Low_Resource_Settings.

Buys R, et al. Validation of consumer-grade digital camera-based human activity evaluation for upper limb exercises and development of a therapist-guided, automated telerehabilitation framework and platform for stroke rehabilitation. arXiv. 2023. Available from: https://arxiv.org/abs/2311.13088.

Fok-Chi-Seng Fok Kow, et al. MIDAS: Multi-sensorial immersive dynamic autonomous system improves motivation of stroke-affected patients for hand rehabilitation. arXiv. 2022. Available from: https://arxiv.org/abs/2203.10536.

Kantan P, et al. A technical framework for musical biofeedback in stroke rehabilitation. arXiv. 2020. Available from: https://arxiv.org/abs/2012.00323.

Ahmad, N. R. (2025). *Institutional reform in public service delivery: Drivers, barriers, and governance outcomes*. *Journal of Humanities and Social Sciences*. https://doi.org/10.52152/jhs8rn12