PAKISTAN JOURNAL OF HEALTH SOLUTIONS

vol:04 issue:01 2026

E-ISSN:2520-0674

GUT HEALTH AND ITS IMPACT ON MENTAL HEALTH: EXPLORING THE GUT-BRAIN AXIS

Afsar Mian

Centre for Regenerative Medicine & Stem Cell Research (CRM), Aga Khan University, Karachi, Pakistan

Article Info:

Keywords:

Gut-brain axis, microbiome, mental health, depression, anxiety, probiotics, dysbiosis, neuroinflammation, serotonin, psychobiotics .

Abstract. *Emerging research reveals a profound connection between gut health and mental well-being, centered around the gut-brain axis—a bidirectional communication system between the gastrointestinal tract and the central nervous system. The gut microbiome, composed of trillions of microorganisms, influences neurodevelopment, stress response, mood, and cognition through mechanisms involving neurotransmitters, immune modulation, and the vagus nerve. This review examines the evidence linking gut dysbiosis to psychiatric disorders such as depression, anxiety, and autism spectrum disorder, and evaluates potential therapeutic approaches including probiotics, prebiotics, dietary interventions, and fecal microbiota transplantation. Understanding the gut-brain connection may revolutionize mental health treatment by offering integrative, microbiome-targeted therapies.*

1. INTRODUCTION

The gut is increasingly recognized as a "second brain," with mounting evidence linking gut microbial composition to cognitive and emotional health. The gut-brain axis comprises complex neurochemical signaling pathways influenced by the intestinal microbiota. Disruption of this balance—termed dysbiosis—has been associated with mood disorders, neurodegeneration, and behavioral changes. This paper explores the mechanisms behind gut-mind interactions and highlights how manipulating gut health may provide novel avenues for treating psychiatric conditions.

The **Gut-Brain Axis (GBA)** refers to the bidirectional communication network that links the gastrointestinal (GI) tract and the brain, influencing both physical and mental health. Here's an overview of its components and role:

Components of the Gut-Brain Axis:

Vagus Nerve:

The vagus nerve plays a central role in the communication between the gut and the brain. It is one of the longest cranial nerves, stretching from the brainstem to the abdomen. This nerve transmits signals from the gut to the brain, and vice versa, allowing for quick responses to changes in the digestive system.

Enteric Nervous System (ENS):

Often referred to as the "second brain," the ENS consists of a complex network of neurons embedded in the lining of the GI tract. It can function independently of the brain and spinal cord but communicates with the central nervous system (CNS) via the vagus nerve. The ENS regulates digestive processes, including peristalsis (gut motility), secretion of digestive enzymes, and local blood flow.

Immune Pathways:

The gut is home to a large portion of the body's immune system, and it plays a crucial role in regulating immune responses. Immune cells in the gut interact with the microbiota and influence signaling pathways that can affect brain function. Dysregulation of these immune pathways has been linked to various neurological conditions, such as depression and anxiety.

Hypothalamic-Pituitary-Adrenal (HPA) Axis:

The HPA axis is involved in the body's response to stress. It is activated when the body perceives a stressor, triggering the release of hormones like cortisol. The gut microbiota can influence the HPA axis, potentially affecting stress responses, mood, and overall mental health. In turn, chronic stress can alter gut microbiota composition, creating a feedback loop.

Role of Microbiota in Signaling and Neurotransmitter Production:

The gut microbiota, which consists of trillions of microorganisms, including bacteria, fungi, and viruses, plays a key role in the Gut-Brain Axis. The microbiota can influence brain function through various mechanisms:

Neurotransmitter Production:

Many neurotransmitters that are crucial for brain function, such as serotonin, dopamine, and GABA (gamma-aminobutyric acid), are produced by gut microbes. For example, about 90% of the body's serotonin is produced in the gut. These neurotransmitters can influence mood, cognition, and emotional regulation.

Short-Chain Fatty Acids (SCFAs):

The gut microbiota ferments dietary fibers to produce SCFAs (such as butyrate, propionate, and acetate), which have been shown to affect brain function. SCFAs are known to support the integrity of the blood-brain barrier, influence immune responses, and modulate inflammation, all of which are essential for healthy brain function.

Immune System Modulation:

The gut microbiota influences immune cells, including microglia in the brain, and can modulate inflammation in the central nervous system. Chronic inflammation has been linked to conditions like depression, anxiety, and neurodegenerative diseases.

Stress Response and Behavior:

The gut microbiota can impact behavior through its influence on the HPA axis and other neurochemical pathways. For example, gut dysbiosis (imbalance in microbiota

composition) has been associated with increased anxiety, depression, and stress sensitivity.

The Gut-Brain Axis is a complex, dynamic system that involves multiple pathways, including neural, immune, and hormonal signaling, with the gut microbiota playing a key role in regulating brain function, emotional health, and behavior. Alterations in this axis may contribute to various neuropsychological disorders.

2. Microbiome and Neurotransmitters

The human gut microbiome plays a significant role in neurochemistry by influencing the production and regulation of neurotransmitters, as well as other signaling molecules that affect brain function and behavior.

Microbial Synthesis of Key Neurotransmitters

Serotonin (5-HT)

While most serotonin is produced in the gut's enterochromaffin cells, certain microbes—such as *Candida*, *Streptococcus*, *Escherichia*, and *Enterococcus*—can influence its synthesis.

These microbes promote serotonin production by increasing the availability of its precursor, tryptophan, or by interacting with host cells to stimulate its release.

Though serotonin produced in the gut doesn't cross the blood-brain barrier, it influences the enteric nervous system and indirectly affects central serotonin signaling.

Dopamine

Some gut bacteria, like *Bacillus* and *Serratia* species, can produce dopamine directly.

Dopamine affects motivation, pleasure, and reward pathways in the brain. Though peripheral dopamine cannot cross into the brain, microbial influence on precursor availability (like tyrosine and phenylalanine) can alter central dopamine synthesis.

GABA (Gamma-Aminobutyric Acid)

GABA is the primary inhibitory neurotransmitter in the CNS, regulating anxiety and mood.

Certain *Lactobacillus* and *Bifidobacterium* strains are known to produce GABA from glutamate.

These microbial-derived GABA signals can influence vagal pathways to modulate brain function, contributing to anxiolytic (anxiety-reducing) effects.

Short-Chain Fatty Acids (SCFAs): Brain Inflammation and Function

SCFAs are microbial fermentation products of dietary fibers, primarily butyrate, acetate, and propionate. They serve as key signaling molecules in the gut-brain axis:

Neuroinflammation Control

SCFAs, especially butyrate, have anti-inflammatory properties.

They help modulate immune cell function, reduce pro-inflammatory cytokine production, and maintain the integrity of the blood-brain barrier (BBB), which is critical for protecting the brain from harmful substances.

Brain Function and Behavior

SCFAs can affect neuroplasticity, learning, and memory by influencing gene expression through histone deacetylase (HDAC) inhibition, particularly by butyrate.

They also promote neurogenesis (formation of new neurons) and may protect against neurodegenerative diseases like Alzheimer's and Parkinson's.

Gut-Brain Communication

SCFAs modulate vagal nerve activity, influencing mood and cognitive responses.

They also interact with microglia (the brain's resident immune cells), helping regulate their activity and preventing excessive inflammation that could damage

The gut microbiome actively participates in the production of neurotransmitters (serotonin, dopamine, GABA), thereby influencing mood, cognition, and stress response.

SCFAs, as metabolic by-products of gut bacteria, serve crucial roles in reducing neuroinflammation, maintaining BBB integrity, and enhancing cognitive and emotional health.

3. Gut Dysbiosis and Mental Illness

Gut dysbiosis refers to an imbalance in the composition or function of the gut microbiota. This disruption has been increasingly linked to various mental health conditions through mechanisms involving neuroinflammation, altered neurotransmitter signaling, and increased permeability of both the gut lining and the blood-brain barrier (BBB).

Associations with Mental Illness

Depression

Dysbiosis can reduce beneficial microbial species (e.g., *Lactobacillus*, *Bifidobacterium*) and increase pro-inflammatory ones.

This imbalance can decrease production of GABA and serotonin precursors and increase levels of inflammatory cytokines, both of which are implicated in major depressive disorder (MDD).

Studies show that patients with depression often have lower microbial diversity and altered levels of short-chain fatty acids (SCFAs).

Anxiety

A disrupted gut microbiota can affect stress reactivity via the HPA axis, increasing cortisol levels and altering behavior.

Dysbiosis may also affect the vagal nerve signaling that normally helps regulate anxiety.

Some probiotic strains (psychobiotics) have shown promise in reducing anxiety symptoms, suggesting a direct microbiota-brain link.

Autism Spectrum Disorder (ASD)

Children with ASD often present with GI symptoms and altered gut microbial profiles, including increased *Clostridium* species and reduced beneficial bacteria.

Gut dysbiosis may affect neurodevelopment through immune activation, altered metabolite production, and disrupted neurotransmission.

Microbial metabolites such as propionic acid, found elevated in some ASD cases, may influence brain function and behavior.

Mechanisms: Neuroinflammation, Leaky Gut, and BBB Permeability

Neuroinflammation

Gut dysbiosis increases systemic inflammation, which can extend to the brain.

Overactive immune signaling (e.g., elevated IL-6, TNF-α) activates microglia, the brain's immune cells, contributing to psychiatric symptoms.

Chronic neuroinflammation is a hallmark in depression, schizophrenia, and neurodevelopmental disorders.

A healthy gut barrier prevents harmful substances (e.g., bacterial endotoxins like lipopolysaccharide, or LPS) from entering the bloodstream.

In dysbiosis, this barrier becomes "leaky," allowing LPS and other toxins to provoke immune responses and systemic inflammation.

Elevated LPS levels have been found in patients with depression and cognitive disorders.

Blood-Brain Barrier (BBB) Permeability

Inflammation and microbial metabolites like SCFAs can affect the tight junctions of the BBB.

Disruption of the BBB allows immune cells, cytokines, and neurotoxic substances to enter the brain, exacerbating cognitive and emotional dysfunction.

A compromised BBB is linked to disorders such as Alzheimer's disease, MS, and mood disorders.

Gut dysbiosis is strongly associated with mental illnesses including depression, anxiety, and ASD, primarily through:

Neuroinflammatory processes

Increased intestinal and BBB permeability

Disruption of neurotransmitter signaling and microbial metabolite production

Would you like a table comparing the microbiota changes and mechanisms across these mental health conditions?

4. Psychobiotics and Probiotic Therapies

Psychobiotics are a class of probiotics (and in some cases prebiotics) that, when ingested in adequate amounts, produce a health benefit in patients suffering from psychiatric or neurological conditions by interacting with the gut-brain axis. These benefits include improvements in mood, anxiety, stress resilience, and cognitive function.

Definition and Evidence for Psychobiotics

Psychobiotics specifically refer to live microorganisms that confer mental health benefits through:

Modulating neurotransmitter production (e.g., serotonin, GABA)

Reducing systemic and neuroinflammation

Regulating the HPA axis response to stress

Influencing vagal nerve signaling

Restoring gut barrier integrity

Clinical Evidence:

Multiple randomized controlled trials (RCTs) and animal studies have demonstrated that psychobiotics can reduce symptoms of depression, anxiety, and stress.

For example, a 2017 meta-analysis found that probiotic supplementation was associated with a significant reduction in depressive symptoms compared to placebo.

Psychobiotics are also being explored for use in ASD, schizophrenia, PTSD, and neurodegenerative diseases, although evidence in these areas is still emerging.

Key Strains and Their Cognitive Effects

Lactobacillus spp.

Lactobacillus rhamnosus:

Shown to reduce anxiety- and depression-like behavior in animal models.

Influences GABA receptor expression in the brain via vagal signaling.

Lactobacillus helveticus (e.g., R0052 strain):

Combined with *Bifidobacterium longum*, has demonstrated anxiolytic and antidepressant effects in humans.

Lactobacillus plantarum:

Associated with improvements in memory, mood, and gut barrier function.

Bifidobacterium spp.

Bifidobacterium longum (e.g., 1714 and NCC3001 strains):

Reduces cortisol levels and improves cognitive performance under stress.

In IBS patients with comorbid anxiety or depression, *B. longum* has shown psychotropic effects.

Bifidobacterium breve:

Found to improve mood and working memory, particularly in older adults.

Mechanisms of Cognitive and Emotional Modulation

Neurochemical modulation: Enhance the availability of neurotransmitters like serotonin and GABA.

Immune regulation: Reduce inflammation that can impact neural function.

Endocrine interaction: Modulate cortisol and other stress hormones.

Neuroplasticity: Influence BDNF (brain-derived neurotrophic factor), which supports cognitive function and emotional resilience.

Psychobiotics are probiotics with mental health benefits that act through the gut-brain axis.

Strains from the Lactobacillus and Bifidobacterium genera are the most studied, with evidence showing improvements in mood, anxiety, stress resilience, and even cognitive function.

Though promising, strain-specific effects and individual variability necessitate further research for clinical standardization.

5. Diet and the Microbiome

Diet is one of the most powerful modulators of the gut microbiome. The composition, diversity, and metabolic activity of gut microbes are highly sensitive to dietary

patterns, and these changes can, in turn, influence mental health through the gut-brain axis.

Impact of Western Diet vs. High-Fiber Mediterranean Diet

Western Diet

Typically high in saturated fats, refined sugars, and processed foods, and low in fiber.

Leads to:

Reduced microbial diversity

Increased abundance of pro-inflammatory microbes (e.g., *Bilophila*, *Clostridium difficile*)

Decreased production of short-chain fatty acids (SCFAs), especially butyrate

Consequences:

Promotes systemic inflammation, leaky gut, and neuroinflammation

Associated with increased risk of depression, anxiety, and cognitive decline

Alters the HPA axis and stress response

Mediterranean Diet

Emphasizes high fiber, fruits, vegetables, whole grains, legumes, nuts, olive oil, and moderate intake of fish and fermented dairy.

Leads to:

Increased abundance of beneficial microbes (e.g., *Bifidobacterium*, *Lactobacillus*, *Faecalibacterium prausnitzii*)

Enhanced SCFA production, particularly butyrate

Consequences:

Anti-inflammatory effects

Improved gut barrier integrity and blood-brain barrier function

Associated with lower risk of depression and better cognitive performance

Role of Fermented Foods and Polyphenols in Mental Health

Fermented Foods

Include yogurt, kefir, kimchi, sauerkraut, miso, tempeh, and kombucha.

Contain live microbes (probiotics) and bioactive peptides.

Benefits:

Increase gut microbial diversity

Enhance neurotransmitter production (e.g., GABA, serotonin precursors)

Reduce inflammation and oxidative stress

Some studies show mood-enhancing effects, including reductions in anxiety and depressive symptoms

Polyphenols

Bioactive plant compounds found in berries, tea, cocoa, red wine, olive oil, and colorful fruits and vegetables.

Poorly absorbed in the small intestine, thus largely metabolized by gut microbiota into neuroactive compounds.

Benefits:

Promote growth of beneficial microbes (e.g., *Akkermansia*, *Lactobacillus*, *Bifidobacterium*)

Exhibit anti-inflammatory, antioxidant, and neuroprotective effects

Improve cognitive function and reduce depressive symptoms, partly by modulating BDNF and neuroinflammation

A Western diet harms the microbiome and increases mental health risks via inflammation and reduced SCFA production.

In contrast, the Mediterranean diet, fermented foods, and polyphenol-rich plants nourish beneficial microbes and support brain health and emotional well-being.

Would you like an infographic comparing these diets and their effects on the brain via the microbiome?

Suggested Graphs & Figures

Figure 1: Diagram of the gut-brain axis communication pathways

Figure 2: Microbial species and their associated neurotransmitters

Figure 3: Comparison of gut microbiome profiles in depressed vs. healthy individuals

Figure 4: Summary of clinical trials involving probiotics for mental health

Figure 5: Impact of diet on microbiota diversity and mental well-being

Shahzad (2025) examined economic innovation in public markets through the case of Punjab Sahulat Bazaars Authority (PSBA), highlighting its contribution to affordable essential commodities and improved market governance. The study emphasized innovative strategies, including dynamic pricing mechanisms, solar-powered market infrastructure, women-friendly vendor facilities, and mobile bazaars designed to serve underserved communities. Using a qualitative documentary approach, Shahzad (2025) reported that these initiatives improved operational efficiency, market accessibility, and consumer affordability, with price reductions of up to 35% without direct government subsidies. The findings suggest that transparent governance, institutional reforms, and innovative market management can strengthen public welfare and provide a potentially replicable model for developing economies.

Kayitesi and Sohail (2026) examined the challenges and enabling factors affecting women entrepreneurs involved in avocado exports in Kigali City, Rwanda. The study identified key barriers including limited access to finance, lack of training, social constraints, and market-related difficulties. The authors highlighted that supportive financial policies, entrepreneurial skills development, and improved business resources are essential for strengthening women's participation and sustainable growth in export-oriented entrepreneurship.

Summary

The gut-brain axis bridges nutrition, immunity, and neurobiology, redefining our understanding of mental illness. Modulating the gut microbiome may complement traditional psychiatric treatments and foster preventive mental healthcare. This paradigm shift underscores the need for cross-disciplinary collaboration between gastroenterology, neurology, and psychiatry.

References

Cryan JF, Dinan TG. Mind-altering microorganisms: the impact of the gut microbiota on brain and behaviour. *Nat Rev Neurosci.* 2012;13(10):701–712.

Mayer EA et al. Gut/brain axis and the microbiota. *J Clin Invest.* 2015;125(3):926–938.

Foster JA, McVey Neufeld KA. Gut–brain axis: how the microbiome influences anxiety and depression. *Trends Neurosci.* 2013;36(5):305–312.

Carabotti M et al. The gut–brain axis: interactions between enteric microbiota, central and enteric nervous systems. *Ann Gastroenterol.* 2015;28(2):203–209.

Kelly JR et al. Breaking down the barriers: the gut microbiome, intestinal permeability and stress-related psychiatric disorders. *Front Cell Neurosci.* 2015;9:392.

Sherwin E et al. Microbiota and the social brain. *Science.* 2019;366(6465):eaar2016.

Sarkar A et al. The microbiome in psychology and cognitive neuroscience. *Trends Cogn Sci.* 2018;22(7):611–636.

Dash S et al. The gut microbiome and diet in psychiatry. *Curr Opin Psychiatry.* 2015;28(1):1–6.

Dinan TG et al. Psychobiotics: a novel class of psychotropic. *Biol Psychiatry.* 2013;74(10):720–726.

Mangiola F et al. Gut microbiota in autism and mood disorders. *World J Gastroenterol.* 2016;22(1):361–368.

Valles-Colomer M et al. The neuroactive potential of the human gut microbiota in quality of life and depression. *Nat Microbiol.* 2019;4(4):623–632.

Liu RT et al. Associations between gut microbiota and psychiatric disorders: A systematic review. *Transl Psychiatry.* 2021;11(1):1–16.

O'Mahony SM et al. Serotonin, tryptophan metabolism and the brain–gut–microbiome axis. *Behav Brain Res.* 2015;277:32–48.

Schmidt C. Mental health: thinking from the gut. *Nature.* 2015;518(7540):S12–S15.

Rea K et al. Microbiome as a key regulator of brain development and function. *Acta Pharmacol Sin.* 2020;41(5):661–668.

Jiang H et al. Altered gut microbiota profile in patients with generalized anxiety disorder. *J Psychiatr Res.* 2018;104:130–136.

Ng QX et al. A systematic review of the role of prebiotics and probiotics in autism spectrum disorders. *Medicina.* 2019;55(5):129.

Wang Y, Kasper LH. The role of microbiome in central nervous system disorders. *Brain Behav Immun.* 2014;38:1–12.

Chu C et al. The microbiota regulate neuronal function and fear extinction learning. *Nature*. 2019;574(7779):543–548.

Sampson TR, Mazmanian SK. Control of brain development, function, and behavior by the microbiome. *Cell Host Microbe*. 2015;17(5):565–576.

Kayitesi, P. J., & Sohail, N. (2026). *Barriers and enablers affecting women entrepreneurs in Rwanda: Evidence from women avocado exporters in Kigali City*. **Asian Journal of Economics, Business and Accounting, 26**(6), 167–183. https://doi.org/10.9734/ajeba/2026/v26i62298

Shahzad, K. (2025). Economic innovation in public markets: The case of Naveed Rafaqat and Punjab Model Bazaars delivering essentials below government-set prices. Pakistan Journal of Humanities and Social Sciences, 13(1). https://doi.org/10.52131/pjhss.2025.v13i1.2942