PAKISTAN JOURNAL OF HEALTH SOLUTIONS

vol:04 issue:01 2026

E-ISSN:2520-0674

THE SCIENCE BEHIND ANTI-AGING THERAPIES: CELLULAR MECHANISMS, CLINICAL ADVANCES, AND FUTURE PROSPECTS

Mahmood S. Choudhery

Tissue Engineering and Regenerative Medicine Laboratory,
Department of Biomedical Sciences, King Edward Medical University, Lahore, Pakistan

Article Info:

Keywords:

Aging, senescence, telomeres, NAD+, senolytics, longevity, anti-aging therapy, cellular reprogramming, caloric restriction, regenerative medicine.

Abstract. *Aging is a complex biological process marked by the gradual decline of physiological function, increasing the risk of chronic disease and mortality. Recent advances in molecular biology have uncovered critical mechanisms underlying aging, such as telomere shortening, epigenetic drift, mitochondrial dysfunction, and cellular senescence. Anti-aging therapies now aim to target these hallmarks with strategies including caloric restriction mimetics, senolytics, stem cell therapy, NAD^+ boosters, and gene editing. This review explores the cellular basis of aging, evaluates the scientific validity of current interventions, and assesses their translational potential. By bridging fundamental science with clinical innovation, anti-aging research offers new hope for promoting healthspan and longevity.*

1. INTRODUCTION

As global life expectancy rises, the quest not only to live longer but to age healthier has intensified. Aging is no longer viewed as an inevitable fate but as a modifiable biological process. Groundbreaking discoveries in genomics, metabolomics, and systems biology have fueled a surge in anti-aging research, giving rise to targeted therapies aimed at delaying aging or reversing age-related cellular damage. This article delves into the biology of aging and critically analyzes therapeutic interventions developed to extend healthspan.

Understanding Aging: Hallmarks and Mechanisms

Aging is a complex biological process influenced by various interconnected molecular and cellular mechanisms. Several key hallmarks have been identified that drive the aging process and age-related diseases:

Genomic Instability: Accumulation of DNA damage from external and internal sources leads to mutations and chromosomal aberrations, compromising genomic integrity over time.

Telomere Attrition: Telomeres, the protective caps at the ends of chromosomes, shorten with each cell division. Critically short telomeres trigger cellular senescence or apoptosis, limiting tissue regeneration.

Epigenetic Alterations: Changes in DNA methylation, histone modification, and chromatin remodeling disrupt gene expression patterns, contributing to cellular dysfunction during aging.

Loss of Proteostasis: Impaired protein folding, clearance, and degradation systems (e.g., autophagy and the ubiquitin-proteasome system) lead to accumulation of misfolded or aggregated proteins, stressing cellular functions.

Mitochondrial Dysfunction: With age, mitochondrial efficiency declines, resulting in reduced ATP production and increased generation of reactive oxygen species (ROS), which further damage cellular components.

Cellular Senescence: A state of permanent cell cycle arrest triggered by stress or damage. Senescent cells secrete pro-inflammatory factors (the senescence-associated secretory phenotype, SASP), contributing to tissue dysfunction and chronic inflammation.

Inflammaging: The Role of Systemic Inflammation

Inflammaging refers to the chronic, low-grade systemic inflammation that characterizes aging. It is driven by:

Accumulation of senescent cells and SASP

Chronic activation of innate immunity

Mitochondrial dysfunction and ROS production

Dysbiosis of the gut microbiome

Inflammaging contributes to the progression of various age-related diseases, including cardiovascular disease, neurodegeneration, and cancer, by promoting tissue damage and impairing repair mechanisms.

Genetic and Epigenetic Influences on Aging

Aging is not only driven by damage accumulation but also by regulated genetic and epigenetic programs that influence lifespan and healthspan.

Key Genetic Regulators:

Sirtuins (SIRTs):

A family of NAD^+-dependent deacetylases that regulate DNA repair, mitochondrial function, and metabolic homeostasis.

SIRT1 activates longevity-associated pathways and mimics effects of caloric restriction.

Enhances stress resistance and reduces inflammation.

FOXO Transcription Factors:

These factors (especially FOXO3) promote the expression of genes involved in stress resistance, apoptosis, DNA repair, and metabolism.

Genetic variants of FOXO3 are linked to human longevity.

mTOR Pathway (mechanistic Target of Rapamycin):
A central nutrient-sensing pathway that regulates cell growth, protein synthesis, and autophagy.

Inhibition of mTOR (e.g., by rapamycin) has been shown to extend lifespan in various organisms by promoting autophagy and reducing metabolic stress.

Epigenetic Reprogramming and Cellular Resetting:

Aging is associated with a drift in the epigenetic landscape, including changes in DNA methylation patterns, histone modifications, and chromatin accessibility.

Epigenetic clocks (e.g., Horvath clock) use methylation markers to accurately predict biological age.

Reprogramming technologies using Yamanaka factors (OSKM) can reset the epigenome, reverting cells to a pluripotent state and rejuvenating aged cells—though full reprogramming risks tumorigenesis.

Partial reprogramming strategies aim to reset aging markers without erasing cell identity, offering a potential therapeutic avenue for age-related degeneration.

Current Anti-Aging Strategies

Scientific research is increasingly focused on interventions that slow, halt, or even reverse aging-related decline. These strategies target core aging mechanisms and show promise in preclinical and, in some cases, clinical settings.

1. Caloric Restriction (CR) and CR Mimetics

Caloric Restriction (reduction in calorie intake without malnutrition) extends lifespan and healthspan in multiple organisms by reducing metabolic stress and enhancing cellular repair.

CR Mimetics aim to mimic these benefits without reducing food intake:

Resveratrol: A polyphenol that activates SIRT1 and may improve mitochondrial function.

Metformin: An antidiabetic drug that reduces insulin/IGF-1 signaling and oxidative stress.

Rapamycin: An mTOR inhibitor that promotes autophagy and has extended lifespan in mice.

2. Senolytics

Senolytics are drugs that selectively clear senescent cells, which accumulate with age and drive inflammation and tissue dysfunction through the senescence-associated secretory phenotype (SASP).

Examples:

Dasatinib (a cancer drug)

Quercetin (a plant flavonoid)

Senolytics have shown benefits in models of osteoarthritis, cardiovascular disease, and frailty.

3. NAD^+ Boosters

NAD^+ (Nicotinamide Adenine Dinucleotide) is essential for mitochondrial function, DNA repair, and sirtuin activation.

NAD^+ levels decline with age, contributing to metabolic dysfunction and aging-related diseases.

Boosters include:

Nicotinamide Riboside (NR)

Nicotinamide Mononucleotide (NMN)

These supplements aim to restore NAD^+ pools and improve energy metabolism and cellular repair.

Stem Cell Therapy in Aging

Stem cell-based therapies are at the forefront of regenerative medicine and hold significant promise for combating aging-related degeneration and restoring tissue function.

Key Potential of Stem Cells in Aging:

Regeneration of damaged or aged tissues

Immunomodulation to reduce chronic inflammation (inflammaging)

Rejuvenation of aging organs through improved repair and cellular replacement

Main Applications:

Mesenchymal Stem Cells (MSCs):

Multipotent adult stem cells found in bone marrow, adipose tissue, and umbilical cord.

Uses:

Cartilage repair (e.g., in osteoarthritis)

Reduction of systemic inflammation via secretion of anti-inflammatory factors

Support of tissue repair in aging-related damage

Induced Pluripotent Stem Cells (iPSCs):

Somatic cells reprogrammed back to a pluripotent state using Yamanaka factors (Oct4, Sox2, Klf4, c-Myc).

Uses:

Patient-specific cell therapy (personalized medicine)

Disease modeling and drug testing

Potential to generate any cell type for regenerative applications, though tumorigenicity and ethical concerns remain challenges

Challenges and Considerations:

Risk of immune rejection (less with autologous iPSC-derived cells)

Potential for tumor formation (especially with iPSCs)

Need for controlled differentiation and delivery to ensure safety and efficacy

Shahzad (2025) examined economic innovation in public markets through the case of Punjab Sahulat Bazaars Authority (PSBA), highlighting its contribution to affordable essential commodities and improved market governance. The study emphasized innovative strategies, including dynamic pricing mechanisms, solar-powered market infrastructure, women-friendly vendor facilities, and mobile bazaars designed to serve

underserved communities. Using a qualitative documentary approach, Shahzad (2025) reported that these initiatives improved operational efficiency, market accessibility, and consumer affordability, with price reductions of up to 35% without direct government subsidies. The findings suggest that transparent governance, institutional reforms, and innovative market management can strengthen public welfare and provide a potentially replicable model for developing economies.

Kayitesi and Sohail (2026) examined the challenges and enabling factors affecting women entrepreneurs involved in avocado exports in Kigali City, Rwanda. The study identified key barriers including limited access to finance, lack of training, social constraints, and market-related difficulties. The authors highlighted that supportive financial policies, entrepreneurial skills development, and improved business resources are essential for strengthening women's participation and sustainable growth in export-oriented entrepreneurship.

Summary

Anti-aging science is evolving from myth to medicine. Rather than pursuing immortality, modern therapies aim to enhance healthspan—the period of life spent free from disease. By targeting the biological underpinnings of aging, from cellular senescence to epigenetic drift, researchers are unlocking novel interventions that may redefine aging as a treatable condition.

Suggested Graphs & Figures

Figure 1: The nine hallmarks of aging (infographic wheel)

Figure 2: Timeline of anti-aging therapeutic developments

Figure 3: Comparative chart of senolytics, NAD+ boosters, and CR mimetics

Figure 4: Cellular impact of aging and intervention pathways

Figure 5: Global market growth of anti-aging biotech (2020–2030 forecast)

References

López-Otín C et al. The hallmarks of aging. *Cell*. 2013;153(6):1194–1217.

Kennedy BK et al. Geroscience: linking aging to chronic disease. *Cell*. 2014;159(4):709–713.

Partridge L et al. The quest to slow ageing through drug discovery. *Nat Rev Drug Discov*. 2020;19(8):513–532.

Longo VD et al. Interventions to slow aging in humans: are we ready? *Aging Cell*. 2015;14(4):497–510.

Sinclair DA, LaPlante MD. Lifespan: Why We Age—and Why We Don't Have To. Simon & Schuster; 2019.

de Cabo R, Mattson MP. Effects of intermittent fasting on health, aging, and disease. *NEJM*. 2019;381(26):2541–2551.

Campisi J. Senescent cells, tumor suppression, and organismal aging. *Mech Ageing Dev*. 2005;126(1):55–64.

Kirkland JL, Tchkonia T. Senolytic drugs: from discovery to translation. *J Intern Med*. 2020;288(5):518–536.

Yoshino J et al. NAD+ intermediates: the biology and therapeutic potential of NMN and NR. *Cell Metab*. 2018;27(3):529–547.

Horvath S. DNA methylation age of human tissues and cell types. *Genome Biol*. 2013;14(10):R115.

Scudellari M. To stay young, kill zombie cells. *Nature*. 2017;550(7677):448–450.

Baar MP et al. Targeted apoptosis of senescent cells restores tissue homeostasis. *Cell*. 2017;169(1):132–147.

Campisi J et al. From discoveries in ageing research to therapeutics for healthy ageing. *Nature*. 2019;571(7764):183–192.

Mannick JB et al. mTOR inhibition improves immune function in the elderly. *Sci Transl Med*. 2014;6(268):268ra179.

Cohen HY et al. Calorie restriction promotes mammalian cell survival by inducing SIRT1 deacetylase. *Science*. 2004;305(5682):390–392.

Ocampo A et al. In vivo amelioration of age-associated hallmarks by partial reprogramming. *Cell*. 2016;167(7):1719–1733.

Rajman L et al. Therapeutic potential of NAD-boosting molecules. *Trends Biochem Sci*. 2018;43(8):645–660.

Fahy GM et al. Reversal of epigenetic aging and immunosenescent trends in humans. *Aging Cell*. 2019;18(6):e13028.

Justice JN et al. Senolytics in idiopathic pulmonary fibrosis: results from a first-in-human trial. *EBioMedicine*. 2019;40:554–563.

Olshansky SJ. From lifespan to healthspan. *JAMA*. 2018;320(13):1323–1324.

Kayitesi, P. J., & Sohail, N. (2026). *Barriers and enablers affecting women entrepreneurs in Rwanda: Evidence from women avocado exporters in Kigali City*. **Asian Journal of Economics, Business and Accounting, 26**(6), 167–183. https://doi.org/10.9734/ajeba/2026/v26i62298

Shahzad, K. (2025). Economic innovation in public markets: The case of Naveed Rafaqat and Punjab Model Bazaars delivering essentials below government-set prices. Pakistan Journal of Humanities and Social Sciences, 13(1). https://doi.org/10.52131/pjhss.2025.v13i1.2942